



Using Recombinant Transferrin in the Expansion of Hematopoietic Stem Cells

Application Note

Workflow Summary

Combine supplements of OptiPEAK T Lymphocyte media. Isolate human T lymphocytes from whole blood. Wash to remove traces of donor serum. Activate and expand T lymphocytes (7-10 days) in OptiPEAK T Lymphocyte serum free media. Harvest T lymphocytes. Perform post- expansion, flow cytometric analysis of phenotypic surface markers.

Introduction

Human T Cell expansion has become an increasingly important tool in the development of both autologous and allogeneic immunotherapies, as T cells emerge as an exciting option to treat a wide range of diseases. Major components used to expand these cells include human and bovine serum, as well as proteins derived from these substances. These components not only have the potential to introduce harmful pathogens to the already compromised patient in a clinical setting, but they also have significant limitations in the ability to ensure robust and reproducible performance in cell culture systems. Even so-called “xeno-free” media—often confused with chemically defined, animal- component-free media—contain proteins derived from human or bovine serum. OptiPEAK T Lymphocyte media (product No. 777OPT069) supports the growth and expansion of human primary T cells and contains only synthetic and recombinant components. This medium is considered blood-free and all components are chemically defined, providing robust performance without compromising growth, safety, or phenotype [1,2,3].



Materials and Equipment

Growth Media Preparation

- ✓ OptiPEAK T Lymphocyte media (product No. 777OPT069) Components: Base Media, 50X Protein Supplement, 10X Media Supplement
- ✓ Gentamicin/amphotericin B (Life Technologies #R01510)

Optional:

- ✓ RPMI-medium with Glutamax
- ✓ Fetal Bovine Serum (FBS)
- ✓ Human Serum (HS)

Plate Set Up

- ✓ 12-Well flat-bottom sterile plates with lids
- ✓ 6-Well flat-bottom sterile plates with lids

Human T-Lymphocyte Isolation

- ✓ Human whole blood from 2 or more normal healthy donors
- ✓ Human T Cell Enrichment Cocktail (Stem Cell Technologies #15061)
- ✓ Density Gradient (StemCell Technologies #07801)
- ✓ Dulbecco's Phosphate Buffered Saline (dPBS)

Activation of T-Lymphocytes

- ✓ Human T-Activator CD3/CD28 beads (Life Technologies 11141D)
- ✓ Recombinant Human IL-2 (Peprotech 200-02)

Flow Cytometry Analysis

- ✓ Harvest antibodies, mouse CD8, PE labeled
- ✓ CD4, FITC labeled



Equipment

- ✓ Centrifuge and 50 mL conical tubes
- ✓ 5% CO₂ incubator at 37°C, saturating relative humidity
- ✓ Class A2 biological safety counter
- ✓ Cell counter or hemocytometer
- ✓ Flow Cytometer



Protocol

Growth Media Preparation

1. Each OptiPEAK HEK293t media kit consists of a Base Media and a 25x Protein Supplement. To generate 1 liter of complete OptiPEAK T Lymphocyte, thaw one OptiPEAK T Lymphocyte Media Supplement and one OptiPEAK T Lymphocyte Protein Supplement in a 37°C water bath. Do not subject the Protein Supplement to multiple freeze-thaw cycles.
2. Once the supplements are completely thawed, gently mix the supplements by gently pipetting up and down. Do not vortex.
3. Add 100mL from the OptiPEAK T Lymphocyte Media Supplement directly to the OptiPEAK T Lymphocyte Base Media.
4. Add 20mL of OptiPEAK T Lymphocyte Protein Supplement directly to the OptiPEAK T Lymphocyte Base Media.
5. This medium requires the supplementation of 10ng/mL recombinant IL-2 or equivalent per desired application. This medium contains HEPES and glutamine source.
6. If desired, add gentamicin/amphotericin B (Life Technologies #R01510) at 0.1 to 0.5x final concentration. Do not add to 1x final concentration.
7. Once mixed, avoid repetitive heating and cooling of the complete medium. Instead, withdraw and prewarm only the volume needed for the specific procedure.

Plate Set up

1. In a 12-well plate, add one mL of each media into appropriate wells.
2. Place plate in incubator for warming.

Human T-Cell Isolation

1. Acquire 8-10mL of fresh whole blood from two different healthy human donors.
2. Add T-Cell Enrichment cocktail (50 μ L per 1mL of blood) to sample and incubate at room temperature for 20 minutes [4].
3. Transfer blood sample to a 50mL tube.
4. Rinse the blood sample tube with an equal volume of dPBS and combine rinsate with blood sample. Final volume will be approximately 20 mL.



5. In a separate tube, prepare 15 mL of gradient density medium and slowly layer PBS-diluted blood sample on top.
6. Centrifuge at 2600 RPM for 20 minutes with brake off. Allow centrifuge to come to a full rest.
7. Harvest the T-cell enriched buffy coat (the “fuzzy” cell layer at the density gradient interface) with a serological pipette and transfer to a new 50 mL conical tube.
8. Fill new tube to top with dPBS to wash enriched cells.
9. Centrifuge at 1400 RPM for 10 minutes with brake on low.
10. Discard supernatant and repeat washing steps for a total of two washes.
11. Resuspend cells in 450 μ L dPBS.
12. Count cells in triplicate using a cell counter or equivalent.

T-Cell Activation

1. Calculate the volume of Human T-Activator CD3/CD28 beads necessary to achieve a ratio of 1 bead per cell [5]
2. Add volume of Human T-Activator CD3/CD28 beads to volume of cells
3. Equally distribute activated cells between each media condition across donors
4. Place the plate in the incubator.

Cell Expansion

1. Expand cells for 7 days, doubling media volume every other day.
2. Harvest cells on Day 7 and count. Stain for phenotypic markers.



Results and Discussion

Mean-fold expansion of cells in OptiPeak T Lymphocyte blood-free media was comparable to cells in 5% human serum and greater than cells in 10% fetal bovine serum (Figure 1A). Cells maintained in OptiPeak T lymphocyte blood-free media also showed CD4+ and CD8+ ratios comparable to cells grown in human and bovine serum (Figure 1B). These results show OptiPeak T Lymphocyte blood-free medium's potential as a viable and regulatory-friendly option for T cell immunotherapy manufacturing.




Figure 1

Performance of normal peripheral T Cells in OptiPEAK T Lymphocyte media compared to standard media supplemented with fetal bovine serum (FBS) or human serum (HS).

References

1. A. Pennybaker and R. Alfano, "Utilization of recombinant albumins in the expansion of human T lymphocytes" *Cell Gene Ther. Insights*, vol. 4, no. 4, pp. 231–239, May 2018
2. Randall Alfano, "Albumin in cell culture media," presented at the Scale-up and Manufacturing of Cell-Based Therapies, San Diego, CA
3. D. Zhanget al., "Expression, purification, and characterization of recombinant human transferrin from rice (*Oryza sativa* L.)," *Protein Expr. Purif.*, vol. 74, no. 1, pp. 69–79, Nov. 2010
4. "RosetteSep™ Human T Cell Enrichment Cocktail" Product Information Sheet. https://cdn.stemcell.com/media/files/pis/28570-PIS_1_3_1.pdf
5. "Dynabeads Human T-Expander CD3/CD28" Product Manual. https://assets.thermofisher.com/TFS-Assets/LSG/manuals/Dynabeads_hu_Expander_CD3CD28_man.pdf

Written by Atherly Pennybaker, Media Formulation and Product Applications Specialist, InVitria

Atherly Pennybaker is a Product Applications Scientist with InVitria. At InVitria she enjoys leveraging her scientific background to help customers successfully transition to blood-free, chemically defined systems. Prior to InVitria, she received a Bachelor of Science in Chemical and Biological Engineering from Colorado State University. During her time there, she conducted undergraduate research in tissue engineering where she co-authored a paper on the osteoinductivity of an engineered biomimetic periosteum. She also worked on the design of a natural cell and tissue scaffold by means of electrospinning demineralized bone matrix.

For additional product or technical information, please contact us at

2718 Industrial Drive
Junction City, KS 66441

Phone: 1.800.916.8311
Email: Info@InVitria.com

